

2014

Utility of Potential Misdiagnoses in Predicting Foodborne Outbreaks

Lucia Lucia

lucia.2009@phdis.smu.edu.sg

Artur Dubrawski

Carnegie Mellon University

Lujie Chen

Carnegie Mellon University

Follow this and additional works at: https://ink.library.smu.edu.sg/sooss_research



Part of the [Social and Behavioral Sciences Commons](#)

Citation

Lucia, Lucia, Dubrawski, Artur, & Chen, Lujie. (2014). Utility of Potential Misdiagnoses in Predicting Foodborne Outbreaks. *Online Journal of Public Health Informatics*, 6(1).

Available at: https://ink.library.smu.edu.sg/sooss_research/1678

This Journal Article is brought to you for free and open access by the School of Social Sciences at Institutional Knowledge at Singapore Management University. It has been accepted for inclusion in Research Collection School of Social Sciences by an authorized administrator of Institutional Knowledge at Singapore Management University. For more information, please email libIR@smu.edu.sg.

Utility of Potential Misdiagnoses in Predicting Foodborne Outbreaks

Lucia Lucia^{*1}, Artur Dubrawski² and Lujie Chen²

¹Singapore Management University, Singapore, Singapore; ²Carnegie Mellon University, Pittsburgh, PA, USA

Objective

To investigate utility of using inpatient and emergency room diagnoses to detect outbreaks of Salmonellosis in humans. To quantify the impact of including in the analysis cases diagnosed with conditions that may have physiological appearance similar to Salmonellosis.

Introduction

Reliable detection and accurate scoping of outbreaks of foodborne illness are the keys to effective mitigation of their impacts. However, relatively small number of persons affected and underreporting, challenge the reliability of surveillance models. In this work, we correlate a record of identified outbreaks and sporadic cases of Salmonellosis in humans retained in PulseNet¹, and diagnosis codes in hospital claims collected in California from 2006 to 2010. We hypothesize that the data support and reliability of detection could be improved by including cases in which Salmonella infection may be confused².

Methods

We join the data in a table indexed with dates and locations, containing counts of inpatient and ED patients diagnosed with Salmonellosis and related diseases, also counts of cases involved in outbreaks, aggregated by day (the admission date or the isolation date) and location (the county of hospital locations or the county where the outbreaks occurred). 9.5% of the 66,845 rows in the table involve sporadic cases and identified clusters.

To quantify predictive utility of potential misdiagnoses, Zero-inflated Poisson regression (ZIP) model³ is trained to predict the number of cases in epidemiological data. Among Salmonellosis (counts in inpatient and ED) and 12 potential misdiagnoses, the best combination of input features is found by exhaustive search to minimize 10 fold cross validation ZIP prediction error. The chosen model is then trained using thusly selected features using all data. Similarly, we train a Random Forest (RF) binary classifier⁴ that also includes spatio-temporal predictors (county and month) to discount seasonality and spatial propensity of outbreaks.

Results

We found that 8 diagnoses related to Salmonellosis have non-trivial impact on outbreak predictability (only Celiac is insignificant with p-value>0.05). Their contributory effect is indicated by positive coefficients of ZIP count model and negative coefficients of ZIP zero model, as shown in the table.

Including counts of these diagnoses improves predictability of the occurrence of outbreaks vs. using Salmonellosis diagnoses only. The AUC score of the RF model increases from 57% to 87%. Adding spatio-temporal factors improves the predictability to 91% AUC. The model discovers 71% of actual outbreak cases at 7% false positive rate (FPr) and correctly recalls 4.5 as many outbreak cases at 1% FPr as when using Salmonellosis diagnoses only.

We found 37% of the predictions can be made 1 to 7 days earlier than the recorded isolation date, increasing precision to 89%. This suggests a potential early warning utility. It is also possible to spot outbreaks not revealed in PulseNet. For instance, 22 out of 35 outbreak predictions in Yolo County are not in PulseNet; 60% of these 22 have

at least 40% of nearby counties showing positive predictions or actual cases in PulseNet in the same periods of time.

Conclusions

Empirically found informative correlation between the counts of hospital patients diagnosed with diseases that may have physiological appearance similar to Salmonellosis, and epidemiologically recorded cases of Salmonellosis. This suggests that tracking these diseases could support accuracy of foodborne illness surveillance. Further study is yet required to verify the actual extent of clinical misdiagnosing, and if there are other factors explaining the apparent correlation.

ZIP Model	Intercept	Salmonellosis emergency	Salmonellosis Inpatient	Shigellosis
Count	-0.477639	0.3227	0.2897	0.1430
Zero	3.47364	-1.1529	-1.0431	-0.0944

ZIP Model	Crohns	Diabetic gastroparesis	Colon cancer	Diarrhea
Count	0.0463	0.0424	0.0348	0.0248
Zero	-0.0445	-0.1459	-0.3631	-0.2986

ZIP Model	Celiac	Ulcerative colitis	Viral gastroenteritis
Count	0.0235	0.0215	-0.0368
Zero	-0.4505	-0.2772	0.0219

Keywords

foodborne outbreaks; misdiagnosis; predictive analytics

Acknowledgments

This work is supported by the National Science Foundation (awards 0911032, 1320347), and the Singapore National Research Foundation under its International Research Centre @Singapore Funding Initiative and administered by the IDM Programme Office, Media Development Authority.

References

1. PulseNet. <http://www.cdc.gov/pulsenet/about/index.html>
2. Rightdiagnosis. <http://www.rightdiagnosis.com>
3. Lambert D. Zero-Inflated Poisson Regression, with an Application to Defects in Manufacturing. *Technometrics*. 1992.34(1): 1-14.
4. Breiman L. Random Forests, *Machine Learning*. 2001.45(1).

*Lucia Lucia

E-mail: lucia.2009@phdis.smu.edu.sg

